

Comprehensive Environmental Assessment of Engineered Aluminum Nanoparticles in DoD Materiel: Evaluation before Acquisition

**Chris Griggs -ERDC -EL
Environmental Engineering Branch**

**Dr. David Johnson-ERDC-EL
Chief, Risk Assessment Branch**

**Dr. Jeff Steevens-ERDC-EL
Project Lead/Senior Scientist**




**US Army Corps of Engineers
BUILDING STRONG®**



Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 28 MAR 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Comprehensive Environmental Assessment of Engineered Aluminum Nanoparticles in DoD Materiel: Evaluation before Acquisition				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Research and Development Center,Environmental Laboratory,3909 Halls Ferry Road,Vicksburg,MS,39180-6199				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the 2011 DoD Environmental Monitoring & Data Quality Workshop (EMDQ 2011), 28 Mar ? 1 Apr, Arlington, VA.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 28	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


Stages of Acquisition Process Benefiting from Environmental Hazard Assessment



Technology development and acquisitions process. Adapted from Mike McDevitt, Installations Management Command




DoD Materiel Development and Costs



- It is estimated that over 85% of the costs of technology occur after systems acquisition



Comprehensive Environmental Assessment (CEA)



Adapted from Davis, 2007




CEA Process

- **Identify the question(s)**
 - ▶ Sources
 - ▶ Life cycle stages, fate & transport, matrices, exposure, effects
 - ▶ Developed methods and standardized protocols
- **Obtain diverse perspectives**
 - ▶ **ODUSD** Chemical & Material Risk Management
 - ▶ **NNCO** National Nanotechnology Coordination Office
 - ▶ **ARMY** - ARDEC , Army Institute of Public Health, ERDC
 - ▶ **Navy** - NSWC-IHD
 - ▶ **Air Force** - Air Force Laboratory Human Effectiveness Directorate
- **Use collective judgment method**

Adapted from Davis, 2007



ERDC CEA Case Study: Engineered Aluminum Nanoparticles




Applying CEA approach to nanotechnology in the R&D Phase

- Lack of mature industries
- Data lacking or evolving
- Characterization of materials
- Uncertainty is high
- Identify and prioritize knowledge gaps



CEA: Life Cycle Stages of nano-AI




Site Visit: ARDEC Picatinny Arsenal Nanotechnology Research Center

- Operates North America's largest Radiofrequency (RF) Induction Plasma reactor (Tekna Plasma Systems) pilot plant for nano-Al and high performance nonmaterial:
 - ▶ Plasma Synthesis
 - ▶ 10 micron aluminum powder feedstock
 - ▶ Use of nano-aluminum still in the R&D phase
 - ▶ Characterization FE-SEM ,XRD, XRF, XRD, BET, Thermal analysis



CEA: Life Cycle Stages – Feedstocks



- Powder feeder system introduces 10 micron aluminum powder
- Plasma system combines “top down” and “bottom up” process, solid precursor vaporized and quenched enabling synthesis of nano-Al
- Feedstock is aluminum oxide passivated in situ with an approximately 5 nm oxide coating
- Nano-scale powder collected in stainless steel canister inside glove box



CEA: Life Cycle Stages – manufacture




Picatinny Arsenal
Nanotechnology Research Center
maintains two 300kiloWatt RF
Tekna Plasma Systems:

- Metals and metal oxides
- Ceramics

- Cyclone classifier separates product by size, nano-scale powder is collected in a collection chamber under argon
- Synthesis efficiency is 10:1 nanopowder to aggregates, aggregates are stored for research purposes



CEA: Life Cycle Stages – manufacture




- HEPA filtered fan with 100 cfm exhaust hood, hydrogen sensors in enclosure to detect leaks
- Reactor process capacity of 4hrs per day with 4 hours remaining to rapidly characterize product FE-SEM ,XRD, XRF, XRD, BET, Thermal Analysis, etc.




CEA: Life Cycle Stages – distribution and storage

- Use of nano-aluminum still in the R&D phase
- Stored under inert atmosphere
- Aggregates are stored at the facility (still have research value),
- Current synthesis of 200g batches for rapid characterization
- Stability studies indicate no loss in surface area, however a 20% loss in reactivity due to oxygen diffusion




CEA: Environmental Pathways of nano-Al




CEA: Environmental Pathways of nano-Al

Technology development and use for nanoaluminum



CEA: Fate and Transport of nano-AI




CEA: Fate and Transport of nano-Al



- Particle size dictates oxidation potential
- Nano-Al/ Al_2O_3 interacts with soil, water, and strongly with humic acids
- Highly agglomerates ➡ affects mobility in soil
- Surface charge changes with leachate ➡ alters mobility
- Micron-sized Al_2O_3 has greater sorption than nano- Al_2O_3



CEA: Exposure-Dose of nano-Al



CEA: Exposure-Dose % content of nano-Al



Plot of extinction values for Al triangular prisms (Faber et al. 2008)



CEA: Exposure-Dose of nano-Al

Most likely routes of
nano-Al/Al₂O₃

exposure: Inhalation > Dermal > Internal (oral, ip, iv)




•**ARDEC-NIOSH collaborative framework** “*Nano-powder Synthesis & Associated Safety Precautions at ARDEC*”

TWA and other occupational exposure values?

R&D laboratory evaluations of occupational exposures?




CEA: Effects of nano-AI




CEA: Problems with Effects of nano-Al

- Nano-Al/ Al_2O_3 is highly agglomerated
- Is aged nano-Al the same as nano- Al_2O_3 ?



Increased Oxidation



CEA: Effects of nano-Al

Ecosystems:

- Less toxic to daphnids and algae than other NPs
- More toxic to juvenile zebrafish than adults
- Causes atherothrombotic events in zebrafish
- Produces differential effects on benthic organisms
- Mildly toxic to bacteria
- Mildly phytotoxic (root growth inhibition) due to ROS
- Soil nematodes and earthworm reproduction negatively affected



CEA: Effects of nano-Al

Human Health:

1. Inhalation –

Negatively affects alveolar macrophages function

2. Dermal –

Dermal contact may increase proinflammation, dermatitis

3. Internal –

Neurotoxicity (blood brain barrier disruption) and

Increased genotoxicity



Preliminary Conclusions

- Potential sources and releases of nano-Al to the environment that will likely occur through air, water, or soil exposures through the production, use, and disposal of nano-Al propellants, igniters, and additives.
- However, these preliminary findings are the result of an assessment from the R&D community.
- Data collection is still required to gain a better understanding of the future deployment and handling of nano-Al as a military technology.




Data Gaps / Moving Forward

- **Life Cycle:** Further collaboration required within the R&D community such ARDEC, NSWC-IHD, and AFRL to discuss life cycle phases.
- **Environmental Pathways:** This is a potential laboratory and field research project (modeling and hyperspectral imaging analysis for nano-Al/energetic combustion analyses)
- **Exposure:** Exposure to biota and humans is perhaps the biggest area of uncertainty in this entire nano-Al CEA.
- **Environmental Fate:** Once nano-Al has moved beyond the R&D phase, field testing will be imperative to study nano-Al/ Al_2O_3 propellants and energetics in the field.
- **Effects:** Data needs to reflect of actual particle sizes, i.e., nanoparticle agglomerates vs. monodispersed nanoparticles.



Questions?



References

Coleman, J.G., Johnson, D.R., Stanley, J.K., Bednar, A.J., Weiss, R.E., Steevens, J.A. (2010) Assessing the fate and effects of nano aluminum oxide in the terrestrial earthworm, *Eisenia fetida*. *Environ. Toxicol. Chem.* 29(7): 1575-1580.

Darlington, T.K., Neigh, A.M., Spencer, M.T., Nguyen, O.T., Oldenburg, S.J. (2009) Nanoparticle characteristics affecting environmental fate and transport through soil. *Environ. Toxicol. Chem.* 28(6): 1191-1199.

Davis, J.M. (2007) How to assess the risks of nanotechnology: learning from past experience. *Nanosci. Nanotech.* 7: 402-409.

Davis, J.M. (2007) How to assess the risks of nanotechnology: learning from past experience. *Nanosci. Nanotech.* 7: 402-409.

Doshi, R., Braid, W., Christodoulatos, C., Wazne, M., O'Connor, G. (2008) Nano-aluminum: transport through sand columns and environmental effects on plants and soil communities. *Environ. Res.* 106: 296-303.

Faber, Benjamin J., Schatz, George C., and Camden, Jon P. *Electrodynamics Calculations of SMSERS Active Junctions and New UV Active Substrates*. *Nanoscope Vol. 5*: 41-51

Johnson, D.R., Methner, M.M., Kennedy, A.J., Steevens, J.A. (2010) Potential for occupational exposure to engineered carbon-based nanomaterials in environmental laboratory studies. *Environ. Health Perspect.* 118(1): 49-54.

Stanley, J.K., Coleman, J.G., Weiss, C.A., Steevens, J.A. (2010) Sediment toxicity and bioaccumulation of nano and micron-sized aluminum oxide. *Environ. Toxicol. Chem.* 29(2): 422-429.

Critical review and advising from Dr. Mike Davis, Senior Science Advisor, U.S. EPA and Dr. Thomas Seager, Professor, University of Arizona

